

# Vector Spaces

Ashwin Bhola

CDS, NYU


Sept 4<sup>th</sup>, 2019

## Visualizations in $\mathbb{R}^2$


Consider 2 vectors  $v$  and  $w$  in  $\mathbb{R}^2$ . Let  $v = (2,2)$  and  $w = (-2,3)$ . Interpret the following sets geometrically. Which of these are a subspaces of  $\mathbb{R}^2$ ?

- $\text{Span}(v)$
- $\text{Span}(v) \cup \text{Span}(w)$
- $\text{Span}(v) \cap \text{Span}(w)$
- $\text{Span}(v, w)$
- $\{(1-t)v + tw : t \in (0,1)\}$
- $\{(1-t)v + tw : t \in \mathbb{R}\}$
- $\{av + bw : a, b \geq 0\}$
- $\{(a, b) \in \mathbb{R}^2 : a^2 + b^2 \leq 25\}$
- $\{(a, a+5) \in \mathbb{R}^2 : a \in \mathbb{R}\}$


# Visualizations in $\mathbb{R}^2$



# Visualizations in $\mathbb{R}^2$




# Visualizations in $\mathbb{R}^2$




$\text{span}(v) \cup \text{span}(w)$

# Visualizations in $\mathbb{R}^2$




$\text{span}(v) \cap \text{span}(w)$

# Visualizations in $\mathbb{R}^2$




# Visualizations in $\mathbb{R}^2$



$span(v, w)$




# Visualizations in $\mathbb{R}^2$




$\text{span}(v, w)$

# Visualizations in $\mathbb{R}^2$




$$tv + (1 - t)w, t \in [0,1]$$

# Visualizations in $\mathbb{R}^2$




$$tv + (1 - t)w, t \in \mathbb{R}$$

# Visualizations in $\mathbb{R}^2$




# Visualizations in $\mathbb{R}^2$




$$av + bw \mid a, b \geq 0$$

# Visualizations in $\mathbb{R}^2$



$$a^2 + b^2 \leq 25$$

# Visualizations in $\mathbb{R}^2$



$$(a, a + 5) \quad a \in \mathbb{R}$$

# Linear Independence, Span, Basis and Dimension

1. Let  $V := \mathbb{R}^{n \times n}$  be the space of  $n \times n$  matrices. Prove that  $V$  is a real vector space. Find the dimension of  $V$ . Let  $U$  be the space of  $n \times n$  diagonal matrices. Is  $U$  a subspace of  $V$ ? What is the dimension of  $U$ ?
2. Let  $v_1, v_2, v_3, v_4$  (all distinct)  $\in \mathbb{R}^3$  and  $C_1 = \{v_1, v_2\}$ ;  $C_2 = \{v_3, v_4\}$ . If  $C_1$  and  $C_2$  are both linearly independent, what are the possible values for  $\dim(\text{Span}(v_1, v_2, v_3, v_4))$ ? No proof necessary
3. True or False: If  $B$  is a basis of  $\mathbb{R}^n$  and  $W$  is a subspace of  $\mathbb{R}^n$ , then a subset of  $B$  is the basis of  $W$
4. Consider the non-empty set of functions  $V := \{p: \mathbb{R} \rightarrow \mathbb{R} \mid p(x) = \sum_{k=0}^n a_k x^k \text{ for } a_k \in \mathbb{R}, \text{ and } x \in \mathbb{R} \text{ is a constant}\}$ . Define an addition operation  $+: V \times V \rightarrow V$  and a scalar multiplication operation  $\cdot: \mathbb{R} \times V \rightarrow V$  such that the triple  $(V, +, \cdot)$  is a real vector space. Find a basis of this vector space and deduce its dimension
5. Suppose  $(v_1, v_2, \dots, v_m) \in \mathbb{R}^n$  be linearly dependent. Prove that for  $x \in \text{span}(v_1, v_2, \dots, v_m)$ , there exist infinitely many  $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_m) \in \mathbb{R}^m$  such that  $x = \sum \alpha_i v_i$